

GI Solutions  
7447 W. Talcott Ave. Suite 209  
Chicago, IL. 60631

Office Phone: 773-631-2728  
Answering Service: 312-492-5764



## **OFFICE BRAVO**

Date of Procedure: \_\_\_\_/\_\_\_\_/\_\_\_\_ Time: \_\_\_\_am/pm  
Report to suite 207, a half hour before your procedure time.  
Please report at: \_\_\_\_am/pm

Nothing to eat or drink after midnight (including NO WATER and NO MEDICATIONS, except for blood pressure and heart medications which may be taken until 7 am the morning of the procedure)

All PPI medications must be stopped one week prior to procedure (ex. Dexilant, Nexium, Omeprazole, Pantoprazole, Tums, Maalox, etc.)

Please inform the office if you have a nickel allergy.

If you are currently taking an aspirin, aspirin products, or blood thinners please see the hand out that has been provided for dosing instructions.

If you are on Coumadin or Warfarin you must have a bleeding profile blood test done the day before or the morning of the procedure.

If you are diabetic and are on insulin contact your primary care physician regarding modification of your dosage.

If you are experiencing severe reflux you can take over the counter Pepcid or famotidine/ranitidine up to two days before the procedure.

Female patients that are not post-menopausal (no period in over one year) are required to give a urine sample before the procedure.

If you have loose teeth they will need to be fixed prior to procedure; failure to do so will result in the cancellation of your procedure and a \$100 fee that is non-refundable through insurance.

Since you will receive IV anesthesia for this procedure you may not drive yourself home. Make sure you make arrangements for a ride home after the gastroscopy.

You will need to schedule an office follow up to review the results approximately one week after your procedure.

Please do not wear jewelry or nail polish/acrylic nails for your procedure(s).